

Multi-resolution in image processing

Preliminaries on wavelets

Barbara Pascal[†]

bpascal-fr.github.io/talks

June 2026

[†] Nantes Université, École Centrale Nantes, CNRS, LS2N, F-44000 Nantes, France



Multiple scales in real data



Multiple scales in real data




Multiple scales in real data



Multiple scales in real data




Multiple scales in real data



Overview of the history of wavelets

Revealing evolutions in dynamical networks

Matteo Morini, Patrick Flandrin, Eric Fleury, Tommaso Venturini, Pablo Jensen¹
IXXI, ENS de Lyon, INRIA, CNRS, LIP UMR 5668, LP UMR 5672




Wavelet Transforms

Continuous Wavelet Transform

Wavelet: $\psi \in L^2(\mathbb{R})$ locally oscillating, integrable with $\int_{\mathbb{R}} \psi(s) ds = 0$

Example: real Shannon wavelet

$$\psi(t) = \frac{\sin(2\pi t) - \sin(\pi t)}{\pi t}$$




Continuous Wavelet Transform


Wavelet: $\psi \in L^2(\mathbb{R})$ locally oscillating, integrable with $\int_{\mathbb{R}} \psi(s) ds = 0$

Example: real Shannon wavelet

$$\psi(t) = \frac{\sin(2\pi t) - \sin(\pi t)}{\pi t}$$



$$\psi_{t,a}(s) = \frac{1}{\sqrt{a}} \psi\left(\frac{s-t}{a}\right)$$




$\implies \{\psi_{t,a}, (t, a) \in \mathbb{R} \times \mathbb{R}_+\}$ atoms with **different** time supports

Continuous Wavelet Transform


Wavelet: $\psi \in L^2(\mathbb{R})$ locally oscillating, integrable with $\int_{\mathbb{R}} \psi(s) ds = 0$

Example: real Shannon wavelet

$$\psi(t) = \frac{\sin(2\pi t) - \sin(\pi t)}{\pi t}$$



$$\psi_{t,a}(s) = \frac{1}{\sqrt{a}} \psi\left(\frac{s-t}{a}\right)$$




$\implies \{\psi_{t,a}, (t, a) \in \mathbb{R} \times \mathbb{R}_+\}$ atoms with **different** time supports

Continuous Wavelet Transform


Wavelet: $\psi \in L^2(\mathbb{R})$ locally oscillating, integrable with $\int_{\mathbb{R}} \psi(s) ds = 0$

Example: real Shannon wavelet

$$\psi(t) = \frac{\sin(2\pi t) - \sin(\pi t)}{\pi t}$$



$$\psi_{t,a}(s) = \frac{1}{\sqrt{a}} \psi\left(\frac{s-t}{a}\right)$$




$\implies \{\psi_{t,a}, (t, a) \in \mathbb{R} \times \mathbb{R}_+\}$ atoms with **different** time supports

Continuous Wavelet Transform

Wavelet: $\psi \in L^2(\mathbb{R})$ locally oscillating, integrable with $\int_{\mathbb{R}} \psi(s) ds = 0$

Example: real Shannon wavelet

$$\psi(t) = \frac{\sin(2\pi t) - \sin(\pi t)}{\pi t}$$



Continuous Wavelet Transform of a finite-energy signal $f \in L^2(\mathbb{R})$

$$\mathcal{W}_f(t, a) = \langle f, \psi_{t,a} \rangle = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{a}} \overline{\psi\left(\frac{s-t}{a}\right)} ds$$

$\langle \cdot, \cdot \rangle$ scalar product in $L^2(\mathbb{R})$, $\bar{\cdot}$ complex conjugate

$\implies \{\psi_{t,a}, (t, a) \in \mathbb{R} \times \mathbb{R}_+\}$ atoms with **different** time supports

Example of a gravitational wave

Physics: two objects $M = m_1 + m_2$ at distance R , $\mu^{-1} = m_1^{-1} + m_2^{-1}$

$$f(t) = A(t_0 - t)^{-1/4} \cos(d(t_0 - t)^{5/8} + \varphi) \mathbf{1}_{(-\infty; t_0[}(t)$$

chirp: amplitude $a(t) = A(t_0 - t)^{-1/4}$, frequency $\omega(t) = \frac{10\pi d}{8}(t_0 - t)^{-3/8}$

– t_0 : time of coalescence,

– d : instantaneous frequency parameter $d \simeq 241 \mathcal{M}_\odot^{-5/8}$,

– A : amplitude reference $A \simeq 3.37 \times 10^{-21} \mathcal{M}_\odot^{5/4} / R$.

Unknown: $\mathcal{M}_\odot = \mu^{3/5} M^{2/5} / M_\odot$: chirp mass in solar mass unit M_\odot

Example of a gravitational wave

Physics: two objects $M = m_1 + m_2$ at distance R , $\mu^{-1} = m_1^{-1} + m_2^{-1}$

$$f(t) = A(t_0 - t)^{-1/4} \cos(d(t_0 - t)^{5/8} + \varphi) \mathbf{1}_{(-\infty; t_0]}(t)$$


chirp: amplitude $a(t) = A(t_0 - t)^{-1/4}$, frequency $\omega(t) = \frac{10\pi d}{8}(t_0 - t)^{-3/8}$

– t_0 : time of coalescence,

– d : instantaneous frequency parameter $d \simeq 241 \mathcal{M}_\odot^{-5/8}$,

– A : amplitude reference $A \simeq 3.37 \times 10^{-21} \mathcal{M}_\odot^{5/4} / R$.

Unknown: $\mathcal{M}_\odot = \mu^{3/5} M^{2/5} / M_\odot$: chirp mass in solar mass unit M_\odot



LIGO Open Science Center

Example of a gravitational wave

Physics: two objects $M = m_1 + m_2$ at distance R , $\mu^{-1} = m_1^{-1} + m_2^{-1}$

$$f(t) = A(t_0 - t)^{-1/4} \cos(d(t_0 - t)^{5/8} + \varphi) \mathbf{1}_{(-\infty; t_0[}(t)$$

chirp: amplitude $a(t) = A(t_0 - t)^{-1/4}$, frequency $\omega(t) = \frac{10\pi d}{8}(t_0 - t)^{-3/8}$

– t_0 : time of coalescence,


– d : instantaneous frequency parameter

$$d \simeq 241 M_{\odot}^{-5/8},$$


– A : amplitude reference

$$A \simeq 3.37 \times 10^{-21} M_{\odot}^{5/4} / R.$$

Unknown: $\mathcal{M}_{\odot} = \mu^{3/5} M^{2/5} / M_{\odot}$: chirp mass in solar mass unit M_{\odot}



LIGO Open Science Center



Morlet scalogram $|\mathcal{W}_f(t, a)|^2$

tftb.nongnu.org

Example of a gravitational wave

Physics: two objects $M = m_1 + m_2$ at distance R , $\mu^{-1} = m_1^{-1} + m_2^{-1}$

$$f(t) = A(t_0 - t)^{-1/4} \cos(d(t_0 - t)^{5/8} + \varphi) \mathbf{1}_{(-\infty; t_0]}(t)$$


chirp: amplitude $a(t) = A(t_0 - t)^{-1/4}$, frequency $\omega(t) = \frac{10\pi d}{8}(t_0 - t)^{-3/8}$

– t_0 : time of coalescence,

– d : instantaneous frequency parameter $d \simeq 241 \mathcal{M}_\odot^{-5/8}$,

– A : amplitude reference $A \simeq 3.37 \times 10^{-21} \mathcal{M}_\odot^{5/4} / R$.

Unknown: $\mathcal{M}_\odot = \mu^{3/5} M^{2/5} / M_\odot$: chirp mass in solar mass unit M_\odot



LIGO Open Science Center

Example of a gravitational wave

Physics: two objects $M = m_1 + m_2$ at distance R , $\mu^{-1} = m_1^{-1} + m_2^{-1}$

$$f(t) = A(t_0 - t)^{-1/4} \cos(d(t_0 - t)^{5/8} + \varphi) \mathbf{1}_{(-\infty; t_0[}(t)$$

chirp: amplitude $a(t) = A(t_0 - t)^{-1/4}$, frequency $\omega(t) = \frac{10\pi d}{8}(t_0 - t)^{-3/8}$

– t_0 : time of coalescence,


– d : instantaneous frequency parameter

$$d \simeq 241 M_{\odot}^{-5/8},$$


– A : amplitude reference

$$A \simeq 3.37 \times 10^{-21} M_{\odot}^{5/4} / R.$$

Unknown: $\mathcal{M}_{\odot} = \mu^{3/5} M^{2/5} / M_{\odot}$: chirp mass in solar mass unit M_{\odot}



LIGO Open Science Center



Morlet scalogram $|\mathcal{W}_f(t, a)|^2$

tftb.nongnu.org

Continuous Wavelet Transform

Reconstruction formula For $\tilde{\psi}$ the Fourier transform of ψ ,

if $C_\psi = \int_{\mathbb{R}} \frac{|\tilde{\psi}(\omega)|^2}{|\omega|} d\omega < \infty$ then ψ is **admissible** and for $f \in L^2(\mathbb{R})$

$$f(t) = \frac{1}{C_\psi} \int_{\mathbb{R} \times \mathbb{R}_+} \mathcal{W}_f(s, a) \frac{1}{\sqrt{a}} \psi\left(\frac{s-t}{a}\right) ds \frac{da}{a^2}$$

with $\mathcal{W}_f(s, a) = \langle f, \psi_{s,a} \rangle$

[A. P. Calderón, 1964, *Stud. Math.*;

A. Grossmann & J. Morlet, 1984, *SIAM J. Math. Anal.*]

Continuous Wavelet Transform

Reconstruction formula For $\tilde{\psi}$ the Fourier transform of ψ ,

if $C_\psi = \int_{\mathbb{R}} \frac{|\tilde{\psi}(\omega)|^2}{|\omega|} d\omega < \infty$ then ψ is **admissible** and for $f \in L^2(\mathbb{R})$

$$f(t) = \frac{1}{C_\psi} \int_{\mathbb{R} \times \mathbb{R}_+} \mathcal{W}_f(s, a) \frac{1}{\sqrt{a}} \psi\left(\frac{s-t}{a}\right) ds \frac{da}{a^2}$$

with $\mathcal{W}_f(s, a) = \langle f, \psi_{s,a} \rangle$ [A. P. Calderón, 1964, *Stud. Math.*;
A. Grossmann & J. Morlet, 1984, *SIAM J. Math. Anal.*]

Reproducing kernel $\mathcal{W}_f(t, a)$ **redundant** representation of f


$$\mathcal{W}_f(t', a') = \frac{1}{C_\psi} \int_{\mathbb{R} \times \mathbb{R}_+} \mathcal{K}(t', t; a', a) \mathcal{W}_f(t, a) dt \frac{da}{a^2}$$

with $\mathcal{K}(t', t; a', a) = \langle \psi_{t,a}, \psi_{t',a'} \rangle$ correlations between wavelets



Continuous Wavelet Transform

Translation invariance


Let $f^{\Delta t}(t) = f(t - \Delta t)$, then $\mathcal{W}_{f^{\Delta t}}(t, a) = \mathcal{W}_f(t - \Delta t, a)$.



$t_0 = 0.44$ [sec]



$t_0 = 0.34$ [sec]



Discrete Signals and Wavelets

From continuous signals to discrete vectors

- f continuous on $[0, 1]$, discretized in $z_n = f\left(\frac{n}{N}\right)$, $n = 0, 1, \dots, N$
 discrete wavelet transform can be computed at scales $N^{-1} < a^j < 1$
- **discrete scales:** $a = 2^{1/v} \implies v$ intermediate scales in octave $[2^j, 2^{j+1})$

Discrete Signals and Wavelets

From continuous signals to discrete vectors

- f continuous on $[0, 1]$, discretized in $z_n = f\left(\frac{n}{N}\right)$, $n = 0, 1, \dots, N$
discrete wavelet transform can be computed at scales $N^{-1} < a^j < 1$
- **discrete scales:** $a = 2^{1/v} \implies v$ intermediate scales in octave $[2^j, 2^{j+1})$

Practical implementation

Integral representation

$$\mathcal{W}_f(t, a) = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{a}} \overline{\psi\left(\frac{s-t}{a}\right)} ds$$

Discrete convolution $z * \overline{\psi_{t,a}(-\cdot)}$

$$\mathcal{W}_z[n, j] = \sum_{m=0}^{N-1} z_m \frac{1}{\sqrt{a^j}} \overline{\psi\left(\frac{m-n}{a^j}\right)}$$

Discrete Signals and Wavelets

From continuous signals to discrete vectors

- f continuous on $[0, 1]$, discretized in $z_n = f\left(\frac{n}{N}\right)$, $n = 0, 1, \dots, N$
discrete wavelet transform can be computed at scales $N^{-1} < a^j < 1$
- **discrete scales**: $a = 2^{1/v} \implies v$ intermediate scales in octave $[2^j, 2^{j+1})$

Practical implementation

Integral representation

$$\mathcal{W}_f(t, a) = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{a}} \overline{\psi\left(\frac{s-t}{a}\right)} ds$$

Discrete convolution $z * \overline{\psi_{t,a}(-\cdot)}$

$$\mathcal{W}_z[n, j] = \sum_{m=0}^{N-1} z_m \frac{1}{\sqrt{a^j}} \overline{\psi\left(\frac{m-n}{a^j}\right)}$$

- **Fast Fourier Transform** at each scale: $\mathcal{O}(N \log_2 N)$ operations

Discrete Signals and Wavelets

From continuous signals to discrete vectors

- f continuous on $[0, 1]$, discretized in $z_n = f\left(\frac{n}{N}\right)$, $n = 0, 1, \dots, N$
discrete wavelet transform can be computed at scales $N^{-1} < a^j < 1$
- **discrete scales**: $a = 2^{1/v} \implies v$ intermediate scales in octave $[2^j, 2^{j+1})$

Practical implementation

Integral representation

$$\mathcal{W}_f(t, a) = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{a}} \overline{\psi\left(\frac{s-t}{a}\right)} ds$$

Discrete convolution $z * \overline{\psi_{t,a}(-\cdot)}$

$$\mathcal{W}_z[n, j] = \sum_{m=0}^{N-1} z_m \frac{1}{\sqrt{a^j}} \overline{\psi\left(\frac{m-n}{a^j}\right)}$$

- **Fast Fourier Transform** at each scale: $\mathcal{O}(N \log_2 N)$ operations
- If ψ with support in $[-K/2, K/2]$: $v \log_2(N/(2K))$ accessible scales

Discrete Signals and Wavelets

From continuous signals to discrete vectors

- f continuous on $[0, 1]$, discretized in $z_n = f\left(\frac{n}{N}\right)$, $n = 0, 1, \dots, N$
discrete wavelet transform can be computed at scales $N^{-1} < a^j < 1$
- **discrete scales**: $a = 2^{1/v} \implies v$ intermediate scales in octave $[2^j, 2^{j+1})$

Practical implementation

Integral representation

$$\mathcal{W}_f(t, a) = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{a}} \overline{\psi\left(\frac{s-t}{a}\right)} ds$$

Discrete convolution $z * \overline{\psi_{t,a}(-\cdot)}$

$$\mathcal{W}_z[n, j] = \sum_{m=0}^{N-1} z_m \frac{1}{\sqrt{a^j}} \overline{\psi\left(\frac{m-n}{a^j}\right)}$$

- **Fast Fourier Transform** at each scale: $\mathcal{O}(N \log_2 N)$ operations
- If ψ with support in $[-K/2, K/2]$: $v \log_2(N/(2K))$ accessible scales


complexity $\mathcal{O}(vN (\log_2 N)^2)$

[S. Mallat, 2009, *Academic Press, Elsevier*]

Multiresolution Analysis

Aims and principles of multiresolution analysis

Motivation: process only details at relevant **discrete** resolutions



[P. J. Burt & E. H. Adelson, 1983, *Proc. IEEE Int. Conf. Commun.*]

Multiresolution analysis

Definition: A multiresolution analysis of $L^2(\mathbb{R})$ is a subspaces sequence

$\{0\} \subset \dots \subset V_1 \subset V_0 \subset V_{-1} \subset \dots \subset V_{-j} \dots \subset V_{-(j+1)} \subset \dots \subset L^2(\mathbb{R})$ satisfying

Multiresolution analysis

Definition: A multiresolution analysis of $L^2(\mathbb{R})$ is a subspaces sequence

$\{0\} \subset \dots \subset V_1 \subset V_0 \subset V_{-1} \subset \dots \subset V_{-j} \dots \subset V_{-(j+1)} \subset \dots \subset L^2(\mathbb{R})$ satisfying

- self-similarity in **time**: $(\forall j \in \mathbb{Z}, f \in V_j, m \in \mathbb{Z}) \quad f(\cdot - m2^j) \in V_j$

Multiresolution analysis

Definition: A multiresolution analysis of $L^2(\mathbb{R})$ is a subspaces sequence

$\{0\} \subset \dots \subset V_1 \subset V_0 \subset V_{-1} \subset \dots \subset V_{-j} \dots \subset V_{-(j+1)} \subset \dots \subset L^2(\mathbb{R})$ satisfying

- self-similarity in **time**: $(\forall j \in \mathbb{Z}, f \in V_j, m \in \mathbb{Z}) \quad f(\cdot - m2^j) \in V_j$
- self-similarity in **scale**: $(\forall j \in \mathbb{Z}) \quad f \in V_j \iff f\left(\frac{\cdot}{2}\right) \in V_{j+1}$

Multiresolution analysis

Definition: A multiresolution analysis of $L^2(\mathbb{R})$ is a subspaces sequence

$\{0\} \subset \dots \subset V_1 \subset V_0 \subset V_{-1} \subset \dots \subset V_{-j} \dots \subset V_{-(j+1)} \subset \dots \subset L^2(\mathbb{R})$ satisfying

- self-similarity in **time**: $(\forall j \in \mathbb{Z}, f \in V_j, m \in \mathbb{Z}) \quad f(\cdot - m2^j) \in V_j$
- self-similarity in **scale**: $(\forall j \in \mathbb{Z}) \quad f \in V_j \iff f\left(\frac{\cdot}{2}\right) \in V_{j+1}$
- **regularity**: φ **father wavelet** or **scaling function** such that
 $\{\varphi(t - k), \quad k \in \mathbb{Z}\}$ is an orthonormal basis of V_0
- **completeness**: $\cup_{j \in \mathbb{Z}} V_j$ is dense in $L^2(\mathbb{R})$ and $\cap_{j \in \mathbb{Z}} V_j = \{0\}$
 $\implies V_j$ **approximation space** at scale 2^j , i.e., *resolution* 2^{-j} .

[P. J. Burt & E. H. Adelson, 1983, *Proc. IEEE Int. Conf. Commun.*;


S. Mallat, 1989, *Trans. Amer. Math. Soc.*;

S. Mallat, 1989, *IEEE Trans. Pattern Anal. Mach. Intell.*;

Y. Meyer, 1992, *Cambridge University Press*]

Aims and principles of multiresolution analysis

Motivation: process only details at relevant **discrete** resolutions



scale 4 \iff resolution 1/4 scale 2 \iff resolution 1/2 scale 1 \iff resolution 1

scale 2^j \iff resolution 2^{-j}

[P. J. Burt & E. H. Adelson, 1983, *Proc. IEEE Int. Conf. Commun.*]

Multiresolution analysis – Daubechies 2 wavelets

From **time** and **scale** invariance and **regularity** condition:

$$\{\sqrt{2^{-j}}\varphi(t/2^j - k), \quad k \in \mathbb{Z}\} \text{ is an orthonormal basis of } V_j$$

Multiresolution analysis – Daubechies 2 wavelets

From **time** and **scale** invariance and **regularity** condition:

$$\{\sqrt{2^{-j}}\varphi(t/2^j - k), \quad k \in \mathbb{Z}\} \text{ is an orthonormal basis of } V_j$$

Orthogonal projection of f onto V_j

$$f|_{V_j} = \sum_{k \in \mathbb{Z}} \phi_{j,k} \sqrt{2^{-j}} \varphi(t/2^j - k)$$

with *approximation coefficients*

$$\phi_{j,k} = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{2^j}} \overline{\varphi(s/2^j - k)} ds$$

Multiresolution analysis – Daubechies 2 wavelets

From **time** and **scale** invariance and **regularity** condition:


$$\{\sqrt{2^{-j}}\varphi(t/2^j - k), \quad k \in \mathbb{Z}\} \text{ is an orthonormal basis of } V_j$$

Orthogonal projection of f onto V_j

$$f|_{V_j} = \sum_{k \in \mathbb{Z}} \phi_{j,k} \sqrt{2^{-j}} \varphi(t/2^j - k)$$

with *approximation coefficients*

$$\phi_{j,k} = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{2^j}} \overline{\varphi(s/2^j - k)} ds$$



Multiresolution analysis – Daubechies 2 wavelets

From **time** and **scale** invariance and **regularity** condition:


$$\{\sqrt{2^{-j}}\varphi(t/2^j - k), \quad k \in \mathbb{Z}\} \text{ is an orthonormal basis of } V_j$$

Orthogonal projection of f onto V_j

$$f|_{V_j} = \sum_{k \in \mathbb{Z}} \phi_{j,k} \sqrt{2^{-j}} \varphi(t/2^j - k)$$

with *approximation coefficients*

$$\phi_{j,k} = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{2^j}} \overline{\varphi(s/2^j - k)} ds$$



Multiresolution analysis – Daubechies 2 wavelets

From **time** and **scale** invariance and **regularity** condition:


$$\{\sqrt{2^{-j}}\varphi(t/2^j - k), \quad k \in \mathbb{Z}\} \text{ is an orthonormal basis of } V_j$$

Orthogonal projection of f onto V_j

$$f|_{V_j} = \sum_{k \in \mathbb{Z}} \phi_{j,k} \sqrt{2^{-j}} \varphi(t/2^j - k)$$

with *approximation coefficients*

$$\phi_{j,k} = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{2^j}} \overline{\varphi(s/2^j - k)} ds$$



Multiresolution analysis – Daubechies 2 wavelets

Approximation and details: $V_{j-1} = V_j \oplus W_j$, W_j : lost information between $f|_{V_{j-1}}$ at resolution $2^{-(j-1)}$ and $f|_{V_j}$ at resolution 2^{-j}

Multiresolution analysis – Daubechies 2 wavelets

Approximation and details: $V_{j-1} = V_j \oplus W_j$, W_j : lost information between $f|_{V_{j-1}}$ at resolution $2^{-(j-1)}$ and $f|_{V_j}$ at resolution 2^{-j}

Theorem There exists a **mother wavelet** $\psi \in L^2(\mathbb{R})$ such that

$\{\psi_{j,k} = \sqrt{2^{-j}}\psi(t/2^j - k), \quad k \in \mathbb{Z}\}$ is an orthonormal basis of W_j .

Multiresolution analysis – Daubechies 2 wavelets

Approximation and details: $V_{j-1} = V_j \oplus W_j$, W_j : lost information between $f|_{V_{j-1}}$ at resolution $2^{-(j-1)}$ and $f|_{V_j}$ at resolution 2^{-j}

Theorem There exists a **mother wavelet** $\psi \in L^2(\mathbb{R})$ such that

$\{\psi_{j,k} = \sqrt{2^{-j}}\psi(t/2^j - k), \quad k \in \mathbb{Z}\}$ is an orthonormal basis of W_j .

Approximation at scale $a = 2^3$



Multiresolution analysis – Daubechies 2 wavelets

Approximation and details: $V_{j-1} = V_j \oplus W_j$, W_j : lost information between $f|_{V_{j-1}}$ at resolution $2^{-(j-1)}$ and $f|_{V_j}$ at resolution 2^{-j}

Theorem There exists a **mother wavelet** $\psi \in L^2(\mathbb{R})$ such that

$\{\psi_{j,k} = \sqrt{2^{-j}}\psi(t/2^j - k), \quad k \in \mathbb{Z}\}$ is an orthonormal basis of W_j .

Orthogonal decomposition of $f|_{V_{j-1}}$

$$f|_{V_{j-1}} = f|_{V_j} + f|_{W_j}$$

with $f|_{W_j} = \sum_{k \in \mathbb{Z}} \zeta_{j,k} \sqrt{2^{-j}} \psi(t/2^j - k)$

involving **wavelet coefficients**

$$\zeta_{j,k} = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{2^j}} \overline{\psi(s/2^j - k)} ds$$

Approximation at scale $a = 2^3$



Multiresolution analysis – Daubechies 2 wavelets

Approximation and details: $V_{j-1} = V_j \oplus W_j$, W_j : lost information between

$f|_{V_{j-1}}$ at resolution $2^{-(j-1)}$ and $f|_{V_j}$ at resolution 2^{-j}

Theorem There exists a **mother wavelet** $\psi \in L^2(\mathbb{R})$ such that

$\{\psi_{j,k} = \sqrt{2^{-j}}\psi(t/2^j - k), \quad k \in \mathbb{Z}\}$ is an orthonormal basis of W_j .

Orthogonal decomposition of $f|_{V_{j-1}}$

$$f|_{V_{j-1}} = f|_{V_j} + f|_{W_j}$$

with $f|_{W_j} = \sum_{k \in \mathbb{Z}} \zeta_{j,k} \sqrt{2^{-j}} \psi(t/2^j - k)$

involving **wavelet coefficients**

$$\zeta_{j,k} = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{2^j}} \overline{\psi(s/2^j - k)} ds$$


Approximation at scale $a = 2^3$



Approximation at scale $a = 2^4$



Details at scale 2^3 disappearing at scale 2^4



Orthonormal wavelet basis

By the **completeness** property of the multiresolution analysis

$$\{\sqrt{2^{-j}}\psi(t/2^j - k), \quad (j, k) \in \mathbb{Z}^2\}$$

is an orthonormal wavelet basis of $L^2(\mathbb{R})$.


Orthonormal wavelet basis

By the **completeness** property of the multiresolution analysis

$$\{\sqrt{2^{-j}}\psi(t/2^j - k), \quad (j, k) \in \mathbb{Z}^2\}$$

is an orthonormal wavelet basis of $L^2(\mathbb{R})$.

Tiling of the time-scale half-plane



Vanishing Moments of Wavelets

Theorem: Let φ scaling function, ψ mother wavelet and $\tilde{\psi}$ its Fourier transform s.t.

$$|\varphi(t)|_{|t| \rightarrow \infty} = \mathcal{O}\left((1+t^2)^{-n_\varphi/2-1}\right), \quad |\psi(t)|_{|t| \rightarrow \infty} = \mathcal{O}\left((1+t^2)^{-n_\psi/2-1}\right)$$

Vanishing Moments of Wavelets

Theorem: Let φ scaling function, ψ mother wavelet and $\tilde{\psi}$ its Fourier transform s.t.

$$|\varphi(t)|_{|t| \rightarrow \infty} = \mathcal{O}((1+t^2)^{-n_\psi/2-1}), \quad |\psi(t)|_{|t| \rightarrow \infty} = \mathcal{O}((1+t^2)^{-n_\psi/2-1})$$

Then the propositions

$$i) \int_{\mathbb{R}} t^n \psi(t) dt = 0, \quad \text{for } n = 0, 1, \dots, n_\psi - 1$$

$$ii) \frac{d^n \tilde{\psi}}{dt^n}(0) = 0, \quad \text{for } n = 0, 1, \dots, n_\psi - 1$$

$$iii) (\forall 0 \leq k < p) \quad t \mapsto \sum_{n=-\infty}^{\infty} n^k \varphi(t-n) \text{ is a polynomial of degree } k \text{ (Fix-Strang)}$$

are equivalent.

[S. Mallat, 2009, *Academic Press, Elsevier*]

Vanishing Moments of Wavelets

Theorem: Let φ scaling function, ψ mother wavelet and $\tilde{\psi}$ its Fourier transform s.t.

$$|\varphi(t)|_{|t| \rightarrow \infty} = \mathcal{O}((1+t^2)^{-n_\psi/2-1}), \quad |\psi(t)|_{|t| \rightarrow \infty} = \mathcal{O}((1+t^2)^{-n_\psi/2-1})$$

Then the propositions

- i) $\int_{\mathbb{R}} t^n \psi(t) dt = 0$, for $n = 0, 1, \dots, n_\psi - 1$
- ii) $\frac{d^n \tilde{\psi}}{dt^n}(0) = 0$, for $n = 0, 1, \dots, n_\psi - 1$
- iii) $(\forall 0 \leq k < p) \quad t \mapsto \sum_{n=-\infty}^{\infty} n^k \varphi(t-n)$ is a polynomial of degree k (Fix-Strang)

are equivalent.

[S. Mallat, 2009, *Academic Press, Elsevier*]

Interpretation:

- mother wavelet **orthogonal** to polynomials of degree at most $n_\psi - 1$
- if signal f is \mathcal{C}^k , $k < n_\psi$

wavelet coefficients $\zeta_{j,n} = \langle f, \psi_{j,n} \rangle$ **small at fine scales**

Support size and number vanishing moments trade-off

If f has an **isolated singularity** at t_0 contained in the support of $\psi_{j,n}$ then the wavelet coefficient $\langle f, \psi_{j,n} \rangle$ is **large**.

Support size and number vanishing moments trade-off

If f has an **isolated singularity** at t_0 contained in the support of $\psi_{j,n}$ then the wavelet coefficient $\langle f, \psi_{j,n} \rangle$ is **large**.

Theorem: If scaling function φ has **compact support** $[a, b]$, then

ψ also has a **compact support** $[(a - b + 1)/2, (b - a + 1)/2]$

of size $b - a$, centered at $1/2$.

Support size and number vanishing moments trade-off

If f has an **isolated singularity** at t_0 contained in the support of $\psi_{j,n}$ then the wavelet coefficient $\langle f, \psi_{j,n} \rangle$ is **large**.

Theorem: If scaling function φ has **compact support** $[a, b]$, then

ψ also has a **compact support** $[(a - b + 1)/2, (b - a + 1)/2]$

of size $b - a$, centered at $1/2$.

If ψ has a compact support of size $\Delta \in \mathbb{N}^*$, at scale 2^j

Δ wavelet coefficients $\zeta_{j,n} = \langle f, \psi_{j,n} \rangle$ are **large**

\implies to reduce the number of significant coefficients, **reduce support size** of ψ .

Support size and number vanishing moments trade-off

Theorem: Let ψ a wavelet with n_ψ vanishing moments generating an orthonormal basis of $L^2(\mathbb{R})$, then its support is of size at least $2n_\psi - 1$.


A **Daubechies** wavelet has a minimum-size support $[-n_\psi + 1, n_\psi]$ and the support of the corresponding scaling function is $[0, 2n_\psi - 1]$.

Support size and number vanishing moments trade-off

Theorem: Let ψ a wavelet with n_ψ vanishing moments generating an orthonormal basis of $L^2(\mathbb{R})$, then its support is of size at least $2n_\psi - 1$.

A **Daubechies** wavelet has a minimum-size support $[-n_\psi + 1, n_\psi]$ and the support of the corresponding scaling function is $[0, 2n_\psi - 1]$.

Haar wavelet: $\psi(t) = \begin{cases} -1 & \text{if } 0 \leq t < 1/2 \\ 1 & \text{if } 1/2 \leq t < 1 \\ \text{otherwise} & \end{cases}$




\Rightarrow **shortest** support among all orthogonal wavelets.

Support size and number vanishing moments trade-off

Theorem: Let ψ a wavelet with n_ψ vanishing moments generating an orthonormal basis of $L^2(\mathbb{R})$, then its support is of size at least $2n_\psi - 1$.


A **Daubechies** wavelet has a minimum-size support $[-n_\psi + 1, n_\psi]$ and the support of the corresponding scaling function is $[0, 2n_\psi - 1]$.

$$\text{Haar wavelet: } \psi(t) = \begin{cases} -1 & \text{if } 0 \leq t < 1/2 \\ 1 & \text{if } 1/2 \leq t < 1 \\ \text{otherwise} & \end{cases}$$




\Rightarrow **shortest** support among all orthogonal wavelets.


Daubechies wavelet:



$n_\psi = 2$



$n_\psi = 3$



$n_\psi = 4$

Decompositions on Wavelet Frames

Theory of Frames

\mathcal{H} : Hilbert space, e.g., $L^2(\mathbb{R})$ or subspace of $L^2(\mathbb{R})$; $\mathbb{I} \subset \mathbb{N}$: set of indices

Definition A family of elements of \mathcal{H} , $\{e_n, n \in \mathbb{I}\}$, s.t.

$$(\forall f \in \mathcal{H}) \quad \underline{\mu} \|f\|^2 \leq \sum_{n \in \mathbb{I}} |\langle f, e_n \rangle|^2 \leq \bar{\mu} \|f\|^2$$

for some bounds $0 < \underline{\mu} \leq \bar{\mu}$, is a **frame**.

Theory of Frames

\mathcal{H} : Hilbert space, e.g., $L^2(\mathbb{R})$ or subspace of $L^2(\mathbb{R})$; $\mathbb{I} \subset \mathbb{N}$: set of indices

Definition A family of elements of \mathcal{H} , $\{e_n, n \in \mathbb{I}\}$, s.t.

$$(\forall f \in \mathcal{H}) \quad \underline{\mu} \|f\|^2 \leq \sum_{n \in \mathbb{I}} |\langle f, e_n \rangle|^2 \leq \bar{\mu} \|f\|^2$$

for some bounds $0 < \underline{\mu} \leq \bar{\mu}$, is a **frame**.

Tight frame If $\underline{\mu} = \bar{\mu} = \mu$, **tight frame** of bound μ .

Union of L orthonormal wavelet bases: **tight frame** of bound L .

Theory of Frames

\mathcal{H} : Hilbert space, e.g., $L^2(\mathbb{R})$ or subspace of $L^2(\mathbb{R})$; $\mathbb{I} \subset \mathbb{N}$: set of indices

Definition A family of elements of \mathcal{H} , $\{e_n, n \in \mathbb{I}\}$, s.t.

$$(\forall f \in \mathcal{H}) \quad \underline{\mu} \|f\|^2 \leq \sum_{n \in \mathbb{Z}} |\langle f, e_n \rangle|^2 \leq \bar{\mu} \|f\|^2$$

for some bounds $0 < \underline{\mu} \leq \bar{\mu}$, is a **frame**.

Tight frame If $\underline{\mu} = \bar{\mu} = \mu$, **tight frame** of bound μ .

Union of L orthonormal wavelet bases: **tight frame** of bound L .

Stable analysis and synthesis

$f \in \mathcal{H} \mapsto (\langle f, e_n \rangle)_{n \in \mathbb{N}} \in \ell^2(\mathbb{I})$ bounded linear operator

$(f_n)_{n \in \mathbb{N}} \in \ell^2(\mathbb{I}) \mapsto \sum_{n \in \mathbb{Z}} f_n e_n \in \mathcal{H}$ bounded linear operator

Theory of Frames

Initially: reconstruction of irregularly sampled band-limited signals

[Duffin & Schaeffer, 1952, *Trans. Amer. Math. Soc.*]

Theory of Frames

Initially: reconstruction of irregularly sampled band-limited signals

[Duffin & Schaeffer, 1952, *Trans. Amer. Math. Soc.*]

Motivations in multiresolution analysis:

Continuous Wavelet Transform: highly redundant, computationally costly

$$\mathcal{W}_f(t, a) = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{a}} \overline{\psi\left(\frac{s-t}{a}\right)} ds$$

Orthonormal wavelet basis decomposition: not invariant under translations

$$(\forall (j, k) \in \mathbb{Z}^2) \quad \zeta_{j,n} = \langle f, \psi_{j,n} \rangle = \sqrt{2^{-j}} \int_{\mathbb{R}} f(t) \overline{\psi(t/2^j - k)} dt$$

Theory of Frames

Initially: reconstruction of irregularly sampled band-limited signals

[Duffin & Schaeffer, 1952, *Trans. Amer. Math. Soc.*]

Motivations in multiresolution analysis:

Continuous Wavelet Transform: highly redundant, computationally costly

$$\mathcal{W}_f(t, a) = \int_{\mathbb{R}} f(s) \frac{1}{\sqrt{a}} \overline{\psi\left(\frac{s-t}{a}\right)} ds$$

Orthonormal wavelet basis decomposition: not invariant under translations

$$(\forall (j, k) \in \mathbb{Z}^2) \quad \zeta_{j,n} = \langle f, \psi_{j,n} \rangle = \sqrt{2^{-j}} \int_{\mathbb{R}} f(t) \psi(t/2^j - k) dt$$

Wavelet frames: $\psi_{j,n}^{(\gamma,b)} = \left\{ \frac{1}{\sqrt{\gamma^j}} \tilde{\psi}\left(\frac{t - bn\gamma^j}{\gamma^j}\right), (j, n) \in \mathbb{Z}^2 \right\}, \gamma > 1, b > 0$

\implies more freedom in the design of the wavelet $\psi^{(\gamma,b)}$

Wavelet Decomposition of Images

Mathematical representation of images



Real-valued square-integrable **field**

$$X : \mathbb{R}^2 \rightarrow \mathbb{R}$$

restricted to a **rectangular** domain $\Omega = [0, n_1 - 1] \times [0, n_2 - 1]$

Two-dimensional wavelet decomposition

Separable wavelet bases:

φ and ψ the scaling function and mother wavelet of a **1D multiresolution analysis**

Define
$$\begin{cases} \psi^{(0)}(\underline{x}) = \varphi(x_1)\varphi(x_2), & \psi^{(1)}(\underline{x}) = \psi(x_1)\varphi(x_2) \\ \psi^{(2)}(\underline{x}) = \varphi(x_1)\psi(x_2), & \psi^{(3)}(\underline{x}) = \psi(x_1)\psi(x_2). \end{cases}$$

Then, the family

$$\left\{ 2^{-j}\psi^{(m)}(\underline{x}2^{-j} - \underline{n}), m \in \{1, 2, 3\}, \underline{x} = (x_1, x_2) \in \mathbb{R}^2, \underline{n} = (n_1, n_2) \in \mathbb{Z}^2 \right\}$$

defines an **orthonormal wavelet basis** of $L^2(\mathbb{R})$.


The **wavelet coefficients** of a 2D field $X \in L^2(\mathbb{R}^2)$ are defined as

$$\zeta_{j,\underline{n}}^{(m)} = \langle X, \psi_{j,\underline{n}}^{(m)} \rangle, \quad \psi_{j,\underline{n}}^{(m)}(\underline{x}) = 2^{-j}\psi^{(m)}(\underline{x}2^{-j} - \underline{n})$$

Wavelet transform of images



Bologna, *Grayscale natural image*




**Daubechies wavelet transform
with $n_\psi = 2$ vanishing moments
at scale 2^1**

Wavelet transform of images



Bologna, *Grayscale natural image*




**Daubechies wavelet transform
with $n_\psi = 2$ vanishing moments
at scale 2^3**

Wavelet transform of images





Bologna, *Grayscale natural image*



Daubechies wavelet transform
with $n_\psi = 2$ vanishing moments
at scale 2^3

Application: compression of images and videos: **JPEG2000, MPEG-4**

References and further readings



Pesquet-Popescu, B., & Pesquet, J. -C. (2001). "Ondelettes et applications", *Techniques de l'ingénieur*, Réf. TE, 5, 215.


Chaux, C. (2006). "Analyse en ondelettes M-bandes en arbre dual; application à la restauration d'images", *Université de Marne la Vallée*.

Pustelnik, N. (2010). "Proximal methods for the resolution of inverse problems: application to positron emission tomography", *Université Paris-Est*.

Multiresolution/multilevel


Multiresolution
to perform
image restoration

(~2000–2015)



Multiresolution
to perform
texture
segmentation

(~2014- now)



Multiresolution
to accelerate
algorithms

(~2016- now)

